

Dollar Spot


[*Sclerotinia homoeocarpa*]

SYMPTOMS

On putting green turf, dollar spot appears as small spots, approximately the size of a dollar coin, that are bleached-white or light tan in color. On turf mowed at heights greater than 0.5", the spots may expand in size up to 6" or more in diameter. The affected leaves typically remain upright and are characterized by having white or light-tan lesions with light reddish-brown margins. As the lesions expand, the leaves are girdled and the upper part of the leaves dies slowly. Distinct lesions are sometimes not evident on close-cut turfgrasses; instead, the leaves die back from the tip and turn white or light tan in color. The grass in the spots may be killed to the soil surface if the disease continues to develop, and many spots may merge to produce large blighted areas. Short, fuzzy white mycelium is often observed on affected turf in the morning when dew is present.



© 2008 Lane Tredway



© 2008 Lane Tredway

dollar spot leaf lesions on
Kentucky bluegrass

Characteristic	Description
Host Grass Species	all
Month(s) with symptoms	February to November
Stand Symptoms	spots, patches (4 to 12 inches)
Foliar Symptoms - Location/Shape	round leaf spots, leaf lesions
Foliar Symptoms - Color	tan or white
Root/Crown Symptoms	none
Fungal Signs	mycelium or none

Note: Still not sure if this is the right disease? The [Turfgrass Disease Identification](#) program may be helpful. Or consult the experts at the [Turf Diagnostics Lab](#). Check the TurfFiles [glossary](#) for definitions of unfamiliar terms.

FACTORS AFFECTING DISEASE DEVELOPMENT

Dollar spot is most common in the spring and fall, when warm, humid days and cool nights lead to heavy dew formation. Dollar spot may develop when night temperatures exceed 50°F and the leaves are wet for 10 to 12 continuous hours. Extended periods of wet, overcast weather can lead to severe dollar spot epidemics on susceptible grasses. The disease can remain active throughout the summer, but activity slows when daytime highs consistently exceed 90°F.

Turfgrasses that are deficient in nutrients, especially nitrogen, are more prone to dollar spot and also recover from the damage more slowly than well-fertilized turf. The disease is also encouraged by drought

stress, low mowing, excessive thatch accumulation, frequent irrigation, and low air movement. Certain cultivars are very susceptible to dollar spot while others are fairly tolerant.

CULTURAL CONTROL

Use of resistant cultivars is one of the most effective means of dollar spot management. This is particularly important for creeping bentgrass and Kentucky bluegrass, as cultivars vary widely in their susceptibility to the disease. Base turfgrass selection on regional cultivar trials and University recommendations, and plant blends and mixtures of cool-season grasses whenever possible.

Adequate nitrogen fertilization will help to prevent dollar spot, and will also encourage plants to recover quickly from the disease if it occurs. Select nitrogen sources, rates, and timings based on local University recommendations for your turfgrass species and climate. Use soil test results to apply the recommended amounts of phosphorus, potassium, lime, and micronutrients.

Dollar spot is encouraged by drought stress and leaf wetness. Proper irrigation timing is needed to balance these factors. Irrigate based on the moisture status of the soil, not on a calendar schedule. Use a soil probe to monitor the amount of moisture in the turfgrass root zone. Irrigate with sufficient water to wet the entire root zone, and then repeat when the entire root zone is no longer moist. Irrigation should be applied early in the morning, between midnight and 6 AM, to keep leaf wetness periods as short as possible. Mowing, dragging, or whipping the turf in the morning to remove dew can help to prevent dollar spot, but these practices can spread the disease if it is actively developing. Improve air movement and reduce humidity by pruning trees, clearing unwanted vegetation, or relocating desirable plants.

Excessive thatch accumulations greatly encourage dollar spot activity. Remove excess thatch by vertical mowing or power raking. Golf course putting greens should be aerified regularly and topdressed with sand to reduce thatch buildup.

Dollar spot is readily spread in leaf tissue or clippings from infected areas. Avoid spreading the disease by washing equipment before entering an uninfected area, by encouraging golfers to clean their shoes between rounds, and by removing and disposing of clippings taken from infected areas.

CHEMICAL CONTROL

Many fungicides control dollar spot, but preventative applications are most effective. A preventative program should be implemented in the early spring when night temperatures consistently exceed 50°F. When applied on a curative basis, fungicides must be applied at high rates and short application intervals.

Uniform spray coverage is important for maximizing fungicide performance; even small gaps in coverage may allow dollar spot to develop. Nozzle type, nozzle pressure, and dilution rate have the greatest impact on the uniformity of fungicide applications. Nozzles that produce coarse to extremely coarse droplets, such as TurfJet or Raindrop nozzles, dramatically reduce the performance of fungicides for dollar spot control. Air-induction or flat fan nozzles that produce fine to medium droplets are recommended. In order to provide thorough coverage of the turfgrass foliage, fungicides should be applied in 2 gallons of water per 1000 ft²; lower carrier volumes reduce the performance of fungicides for foliar disease control.

The fungus that causes dollar spot develops resistance to fungicides very quickly. To prevent or delay the onset of fungicide resistance, use integrated management to minimize fungicide use, rotate among fungicide classes after each application, and tank-mix systemic fungicides with a contact fungicide.

Fungicide	Efficacy ⁽¹⁾	Resistance Risk ⁽²⁾	Class ⁽³⁾	Products ⁽⁴⁾
thiophanate-methyl	++++	9	benzimidazole	3336, Fungo, Systec, T-Storm
boscalid**	++++	9	carboxamide	Emerald

Fungicide	Efficacy ⁽¹⁾	Resistance Risk ⁽²⁾	Class ⁽³⁾	Products ⁽⁴⁾
flutolanil + thiophanate-methyl	++++	9	carboxamide + benzimidazole	SysStar
vinclozolin**	++++	6	dicarboxamide	Curalan, Touche
iprodione + thiophanate-methyl**	++++	9	dicarboxamide + benzimidazole	26/36, Fluid Fungicide, Pro Turf
metconazole	++++	6	DMI	Tourney
myclobutanil	++++	6	DMI	Eagle
propiconazole	++++	6	DMI	Banner MAXX, Propiconazole G-Pro, Propiconazole Pro, Quali-Pro Propiconazole, Savvi, Spectator
triadimefon	++++	6	DMI	Bayleton
triticonazole	++++	6	DMI	Trinity, Triton
triadimefon + trifloxystrobin	++++	9	DMI + QoI	Tartan, Armada
chlorothalonil + thiophanate-methyl**	++++	6	nitrile + benzimidazole	Spectro, ConSyst
chlorothalonil + propiconazole**	++++	4	nitrile + DMI	Concert
chlorothalonil + propiconazole + fludioxonil**	++++	4	nitrile + DMI + phenylpyrolle	Instrata
azoxystrobin + propiconazole	++++	6	QoI + DMI	Headway
iprodione**	+++	6	dicarboxamide	26GT, Iprodione Pro
fenarimol**	+++	6	DMI	Rubigan
chlorothalonil**	+++	3	nitrile	Daconil, Chlorostar, Echo, Manicure
fluoxastrobin	++	9	QoI	Disarm
pyraclostrobin	++	9	QoI	Insignia
mancozeb**	+	3	dithiocarbamate	Fore, Dithane, Pentathlon, Protect
mancozeb + copper hydroxide**	+	3	dithiocarbamate	Junction
thiram**	+	3	dithiocarbamate	Spotrete

** Not for application to residential lawns.

Footnotes:

(1) **Efficacy Codes:**

- ++++ excellent control when conditions are highly favorable for disease development
- +++ good control when disease pressure is high, or excellent control when disease pressure is moderate
- ++ good control when disease pressure is moderate, excellent control when disease pressure is low
- + good control when disease pressure is low
- 0 does not provide adequate control under any conditions
- ? cannot be rated due to insufficient data

(2) Resistance Risk:

- 1 Rotating and tank-mixing not necessary, but recommended to avoid potential side effects from continuous use of same chemical class.
- 2 Rotate to different chemical class after 3-4 applications; tank-mixing not necessary.
- 3 Rotate to different chemical class after 2-3 applications; tank-mixing not necessary.
- 4 Rotate to different chemical class after 1-2 applications; tank-mixing not necessary.
- 6 Rotate to different chemical class after 1-2 applications; tank-mixing with low or moderate risk product recommended.
- 9 Rotate to different chemical class after EVERY application; tank-mix with low or moderate risk product for EVERY application.

(3) Continual use of fungicides with similar control mechanisms (modes of action) can result in fungi that are resistant to some chemicals. Poor or ineffective disease control can be expected when this occurs. Managers can reduce the chances of this happening by mixing or alternating fungicides belonging to different chemical classes.

(4) Recommendations of specific chemicals are based upon information on the manufacturer's label and performance in a limited number of trials. Because environmental conditions and methods of application may vary widely, performance of the chemical will not always conform to the safety and pest control standards indicated by experimental data. When more than one brand name exists for an agricultural chemical, the name of brand that first came onto the market is listed first. Otherwise, brand names are listed in alphabetical order. The order in which brand names are given is not an indication of a recommendation or criticism.

Recommendations for the use of agricultural chemicals are included in this publication as a convenience to the reader. The use of brand names and any mention or listing of commercial products or services does not imply endorsement by North Carolina State University or discrimination against similar products or services not mentioned. Other brand names may be labeled for use on turfgrasses. Individuals who use agricultural chemicals are responsible for ensuring that the intended use complies with current regulations and conforms to the product label. Be sure to obtain current information about usage regulations and examine a current product label before applying any chemical. For assistance, contact your county's Cooperative Extension agent.

Links contained in this document:

Glossary: <http://www.turffiles.ncsu.edu/Glossary.aspx>

Turfgrass Disease Identification Program: <http://www.turffiles.ncsu.edu/diseaseID/>

Turf Diagnostics Lab: <http://ncstateturfdiagnostics.com/TDL/Home.html>

© North Carolina State University. This information sheet was prepared by Lane P. Tredway, Gail G. Wilkerson, Bridget R. Lassiter, Jenifer J. Reynolds, and Gregory S. Buol. Departments of Plant Pathology and Crop Science, College of Agriculture & Life Sciences, North Carolina State University. Prepared March 23, 2009. Available on-line at www.turffiles.ncsu.edu. This publication was made possible through a grant provided by the Center for Turfgrass Environmental Research & Education (CENTERE) whose purpose is to support worthwhile projects that will benefit both the private sector and the public, and protect the environment.